

Computer Science Department

TECHNICAL REPORT

A Property of Graphs of Polytopes

N. Prabhu

Technical Report 454


June 1989

NEW YORK UNIVERSITY



Department of Computer Science
Courant Institute of Mathematical Sciences
251 MERCER STREET, NEW YORK, N.Y. 10012

NYU COMPSCI TR-454
Prabhu, Nagabhushana
A property of graphs of
polytopes.
c.2



A Property of Graphs of Polytopes

N. Prabhu

Technical Report 454

June 1989

A Property of Graphs of Polytopes

Nagabhushana Prabhu*

Courant Institute of Mathematical Sciences
251 Mercer Street
New York, NY 10012

Submitted on June 22, 1989

Abstract

We prove that the subgraph obtained by removing the vertices of a k -face from the graph of a d -polytope ($0 \leq k < d$) is $(d - k - 1)$ -connected. Further we show that this lower bound is tight for $k < d - 1$. We also show that for $k = d - 1$ the known lower bound is tight.

1 Introduction

The 1-skeleton of a polytope P is called the *graph of P* and is denoted $G(P)$. A celebrated result of Balinski shows that *the graph of every d -polytope is d -connected* [1]. The central idea in Balinski's proof is that, *if at a vertex v , a linear functional does not attain the maximum of all its values in the polytope, then v must be adjacent to a vertex at which the linear functional has a higher value.* Using this fact one can easily show that *removing the vertices of a proper face does not disconnect the graph of the polytope* [2]. This corollary provides the background for our discussion.

We address the problem of determining the best lower bound on the connectivity of the remaining subgraph when the vertices of a proper face are removed from the graph of a polytope. In this paper we settle the problem completely. In section 2 we prove the following theorem.

Theorem 1 : *Let P be a d -polytope and Z a k -face of P , $0 \leq k \leq d - 1$. Let $G(P)$ and $G(Z)$ be the graphs of P and Z respectively. Then the complement of $G(Z)$ (i.e. the subgraph of $G(P)$ induced by the vertices that are not in Z) is $(d - k - 1)$ -connected.*

Furthermore, we show that this is a tight lower bound for d -polytopes, when $k < d - 1$. That is, for every $d \in \mathcal{N}$ we can construct a d -polytope which has for every $0 \leq k < d - 1$, a k -face Z such that the complement of $G(Z)$ is *not* $(d - k)$ -connected. When $k = d - 1$ however the aforementioned corollary of Balinski's proof suggests a better lower bound which we prove is tight. That is, for every $d \in \mathcal{N}$ we can construct a d -polytope which contains a facet F such that the complement of $G(F)$ is *not* 2-connected.

*This research was supported in part by NSA grant MDA 904-89-H-2030, ONR grant N00014-85-K-0046 and by NSF grants CCR-8901484, CCR-8902221, CCR-8906949 and CCR-8702271.

2 Connectivity of a subgraph induced by a coface

Let Γ be an abstract graph. Then $V(\Gamma)$ denotes the vertex set of Γ and $E(\Gamma)$ denotes its edge set. Let $V' \subseteq V(\Gamma)$. Let Γ' be the subgraph of Γ induced by V' . By the *complement* of Γ' , denoted $\Gamma \setminus \Gamma'$, we mean the subgraph of Γ induced by the vertex set $V(\Gamma) \setminus V(\Gamma')$.

Let P be a d -polytope and F a proper face of P . Then $V(P)$ and $V(F)$ are the vertex sets of P and F respectively. $G(P)$ is the graph (1-skeleton) of P and $G(F)$ the subgraph of $G(P)$ induced by the vertex set $V(F)$. A subset $C \subset V(P)$ is called a *coface* of P if $F = \text{conv}(V(P) \setminus C)$ is a face of P .

We use the following three results to prove theorem 1.

Result 2.1 ([1,2]) *If M is a proper face of a polytope Q , then the complement of $G(M)$ (i.e. $G(Q) \setminus G(M)$) is connected.*

Result 2.2 ([3]) *A graph $G = (V, E)$ with $|V| \geq k + 1$ is k -connected, if and only if it satisfies the following equivalent conditions.*

1. *If we remove any $k - 1$ vertices from V the subgraph induced by the remaining vertices is still connected.*
2. *Between any two vertices in G there are at least k vertex-disjoint paths.*

Result 2.3 ([3]) *Let $G = (V, E)$ be a graph and let $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ be two subgraphs of G that are k -connected. In addition let V_1 and V_2 have at least k vertices in common. Then the union of G_1 and G_2*

$$G_1 \cup G_2 = (V_1 \cup V_2, E_1 \cup E_2)$$

is also k -connected.

Theorem 1 *Let P be a d -polytope and Z a k -face of P , $0 \leq k \leq d - 1$. Then the complement of $G(Z)$ (i.e. $G(P) \setminus G(Z)$) is $(d - k - 1)$ -connected.*

Proof : The proof of the theorem is by induction on the dimension d . The theorem is trivial when $d = 1$ or $d = 2$. Consider a $d > 2$ and assume that the theorem is true for all n -polytopes, when $1 \leq n < d$. Let P be a d -polytope and Z a k -face of P , $0 \leq k \leq d - 1$. If $k = d - 1$ there is nothing to prove. If $k = d - 2$ we complete the proof at once by appealing to result 2.1. So we assume that $0 \leq k < d - 2$.

Let

$$\mathcal{F} = \{X \mid X \text{ is a facet of } P\}.$$

We partition the set \mathcal{F} into 2 classes \mathcal{A} and \mathcal{B} as follows:

$$\mathcal{A} = \{Y \mid Y \in \mathcal{F}; Y \text{ contains } Z\}$$

$$\mathcal{B} = \{W \mid W \in \mathcal{F}; W \text{ does not contain } Z\}.$$

Note that a facet in \mathcal{B} could have a non-empty intersection with Z . Also observe that both \mathcal{A} and \mathcal{B} are non-empty.

Let $\mathcal{C} \subseteq \mathcal{F}$. Then

$$V(\mathcal{C}) = \bigcup_{F \in \mathcal{C}} V(F).$$

$G(\mathcal{C})$ and $G_Z(\mathcal{C})$ denote the subgraphs of $G(P)$ induced by the vertex sets $V(\mathcal{C})$ and $V(\mathcal{C}) \setminus V(Z)$ respectively.

Consider a facet $X \in \mathcal{B}$. Let $T = X \cap Z$. T is a face of X and $\dim(T) \leq k - 1$. Therefore [4], X contains a $(d - k - 1)$ -face that does not intersect T (and hence disjoint from Z). This shows that $G_Z(X)$ (and hence $G_Z(\mathcal{B})$ and $G_Z(\mathcal{F})$) has at least $d - k$ vertices; so it is meaningful to consider $(d - k - 1)$ -connectedness of $G_Z(\mathcal{B})$ and $G_Z(\mathcal{F})$ (lemmas 2.3 and 2.8).

We first show that $G_Z(\mathcal{B})$ is $(d - k - 1)$ -connected. To do so we need the following notion of ‘connectivity’ of facets. A subset $\mathcal{C} \subseteq \mathcal{F}$ is said to form a ‘*connected*’ complex if for every $X, Y \in \mathcal{C}$ there exists a sequence of facets

$$X = S_1, \dots, S_m = Y$$

such that

- i) $S_j \in \mathcal{C}$, for $1 \leq j \leq m$ and
- ii) for $1 \leq j \leq m - 1$, $S_j \cap S_{j+1}$ is a $(d - 2)$ -face of P .

To distinguish this notion of ‘connectivity’ from *edge connectivity*, in addition to appealing to the context we use quotes in the former case.

Lemma 2.1 \mathcal{B} is a ‘*connected*’ complex.

Proof : Consider a dual P^* of P . Let the k -face Z of P correspond to the $(d - k - 1)$ -face Z^* in P^* . Let X be some facet in \mathcal{B} and X^* the corresponding vertex in P^* . Since X does not contain Z , X^* is not contained in Z^* . Thus the class \mathcal{B} in P corresponds to the set \mathcal{B}^* of all the vertices in P^* that are not contained in Z^* . Since Z^* is a proper face of P^* , using result 2.1 we conclude that the subgraph of $G(P^*)$ induced by \mathcal{B}^* is connected. Therefore \mathcal{B} is a ‘connected’ complex. \bigcirc

Lemma 2.2 For every facet $X \in \mathcal{B}$, $G_Z(X)$ is $(d - k - 1)$ -connected.

Proof : Let $X \in \mathcal{B}$. Let

$$M = X \cap Z.$$

Since X does not contain Z and since the intersection of any two faces of P is again a face of P , M is either a proper face of Z or M is an empty face. In either case $\dim(M) \leq k - 1$. Since X is a $(d - 1)$ -polytope, from the inductive hypothesis we know that the subgraph of $G(X)$ induced by $V(X) \setminus V(M)$ (i.e. $G_Z(X)$) is at least $(d - k - 1)$ -connected. \bigcirc

Lemma 2.3 $G_Z(\mathcal{B})$ is $(d - k - 1)$ -connected.

Proof : Let $|\mathcal{B}| = n$ be the number of facets in \mathcal{B} . Given a subset $\mathcal{S} \subset \mathcal{B}$ which has fewer than n facets, such that $G_Z(\mathcal{S})$ is $(d - k - 1)$ -connected, we show that we can add one more facet $X \in \mathcal{B} \setminus \mathcal{S}$, to \mathcal{S} such that $G_Z(\mathcal{S} \cup \{X\})$ is $(d - k - 1)$ -connected. So if we start with \mathcal{S} containing a single facet from \mathcal{B} , knowing from the preceding lemma that $G_Z(\mathcal{S})$ is $(d - k - 1)$ -connected to start with, we can add facets to \mathcal{S} one at a time in the aforementioned way until $\mathcal{S} = \mathcal{B}$ thus showing $G_Z(\mathcal{B})$ is $(d - k - 1)$ -connected.

Consider an $\mathcal{S} \subset \mathcal{B}$, such that $G_Z(\mathcal{S})$ is $(d - k - 1)$ -connected and $1 \leq |\mathcal{S}| < n$. Since \mathcal{B} is a ‘connected’ complex, there is an $X \in \mathcal{B} \setminus \mathcal{S}$ that shares a $(d - 2)$ -face with some $Y \in \mathcal{S}$. Let

$$W = X \cap Y$$

$$R = X \cap Y \cap Z.$$

R and W are faces of P . Since neither X nor Y contains the k -face Z , $\dim(R) \leq k - 1$. Therefore [4], W contains a face T such that

$$T \cap R = \emptyset$$

and

$$\dim(T) = d - 2 - \dim(R) - 1 \geq d - k - 2.$$

So, $G_Z(Y)$ and $G_Z(X)$ (and hence $G_Z(\mathcal{S})$ and $G_Z(X)$) have at least $d - k - 1$ vertices in common. Moreover since both the subgraphs $G_Z(\mathcal{S})$ and $G_Z(X)$ of $G(P)$ are $(d - k - 1)$ -connected, it follows from result 2.3 that their union is $(d - k - 1)$ -connected. But

$$V(G_Z(\mathcal{S}) \cup G_Z(X)) = V(G_Z(\mathcal{S} \cup \{X\}))$$

and

$$E(G_Z(\mathcal{S}) \cup G_Z(X)) \subseteq E(G_Z(\mathcal{S} \cup \{X\})).$$

Therefore it follows that $G_Z(\mathcal{S} \cup \{X\})$ is $(d - k - 1)$ -connected; that completes the proof. \circ

Before turning to set \mathcal{A} , it is convenient to prove the following lemma.

Lemma 2.4 *Let v be a vertex of a d -polytope Q and let F be some proper face of Q that contains v . Then v has at least one neighbouring vertex in P that does not belong to F .*

Proof : A vertex of a d -polytope together with all its neighbours affinely spans the entire space, E^d [3]. Hence a proper face cannot contain all the neighbours of a vertex in it. \circ

Lemma 2.5 *Every facet in \mathcal{A} is ‘adjacent’ to at least one facet in \mathcal{B} along a $(d - 2)$ -face.*

Proof : The k -face Z of P corresponds to the $(d - k - 1)$ -face Z^* of P^* ; the set \mathcal{A} corresponds to the vertex set $V(Z^*)$ of Z^* and the set \mathcal{B} corresponds to the set of all those vertices of P^* that are not in $V(Z^*)$. Since Z^* is a proper face of P^* , from the previous lemma it follows that every vertex in Z^* is adjacent to at least one vertex that is not in Z^* . \circ

Lemma 2.6 *Let $X \in \mathcal{A}$. Then $G_Z(X)$ and $G_Z(\mathcal{B})$ have at least $d - k - 1$ vertices in common.*

Proof : From the preceding lemma we know that there is a facet Y in \mathcal{B} such that X and Y share a $(d-2)$ -face T , in P . T (a face of Y) shares a face of dimension at most $k-1$, with Z . Therefore T contains a $(d-k-2)$ -face that does not intersect Z . This means T has at least $d-k-1$ vertices none of which is a vertex of Z . So $G_Z(X)$ and $G_Z(Y)$ (and hence $G_Z(X)$ and $G_Z(\mathcal{B})$) share at least $d-k-1$ vertices. \bigcirc

Lemma 2.7 *Let X be a facet in \mathcal{A} . Let $v, w \in V(X) \setminus V(Z)$ be any two vertices of X . Then there are at least $d-k-1$ vertex-disjoint paths between v and w , in $G_Z(\mathcal{F})$.*

Proof : X is a $(d-1)$ -polytope and Z is a k -face of X . From the inductive hypothesis, we conclude that $G_Z(X)$ is $(d-k-2)$ -connected. Therefore there are $d-k-2$ vertex-disjoint paths between v and w , in $G_Z(X)$. Let $n(v), n(w) \in V(P) \setminus V(X)$ be neighbours of v and w respectively (these exist by lemma 2.4). From result 2.1 we know that there exists an edge path Π between $n(v)$ and $n(w)$ that does not pass through any vertex in $V(X)$. Since Z is contained in X and since Π misses X , Π is a path in $G_Z(\mathcal{F})$. The path

$$v \leftrightarrow n(v) \xrightarrow{\Pi} n(w) \leftrightarrow w$$

is vertex-disjoint from every path between v and w in $G_Z(X)$. Together with the $d-k-2$ paths in $G_Z(X)$ we have $d-k-1$ vertex-disjoint paths in all, between v and w . \bigcirc

The following lemma completes the proof of the theorem.

Lemma 2.8 *$G_Z(\mathcal{F})$ is $(d-k-1)$ -connected.*

Proof : Suppose $G_Z(\mathcal{F})$ is not $(d-k-1)$ -connected. Then we can remove a set of $d-k-2$ vertices, say v_1, \dots, v_{d-k-2} , from $G_Z(\mathcal{F})$ such that the remaining subgraph is not connected. Let

$$W = \{v_1, \dots, v_{d-k-2}\}.$$

If $\mathcal{C} \subseteq \mathcal{F}$, then $G_{Z,W}(\mathcal{C})$ denotes the subgraph of $G(P)$ induced by the vertex set $(V(\mathcal{C}) \setminus V(Z)) \setminus W$.

Let C_1, \dots, C_r ($r \geq 2$) be the connected components of $G_{Z,W}(\mathcal{F})$. Since $G_Z(\mathcal{B})$ is $(d-k-1)$ -connected (lemma 2.3) $G_{Z,W}(\mathcal{B})$ is connected; so $G_{Z,W}(\mathcal{B})$ is contained in some connected component, say C_1 . Consider a component C_2 that is different from C_1 . Let v be a vertex in C_2 . Since $v \notin V(C_1)$, $v \notin V(\mathcal{B})$. So $v \in V(\mathcal{A}) \setminus V(\mathcal{B})$. Since $v \in V(\mathcal{A})$, v is a vertex of a facet $Y \in \mathcal{A}$. Lemma 2.6 asserts that $G_Z(Y)$ and $G_Z(\mathcal{B})$ have at least $d-k-1$ vertices in common. Let

$$U = \{m_1, \dots, m_q\} \quad q \geq d-k-1$$

be the set of vertices shared by $G_Z(Y)$ and $G_Z(\mathcal{B})$. Since we removed only $d-k-2$ vertices (when we removed the set W from $G_Z(\mathcal{F})$) at least one vertex of U , say z , remains in $G_{Z,W}(\mathcal{F})$. Both v and z are contained in the facet $Y \in \mathcal{A}$. By appealing to lemma 2.7 we conclude that there are at least $d-k-1$ vertex-disjoint paths between v and z , in $G_Z(\mathcal{F})$. Removing the $d-k-2$ vertices of W from $G_Z(\mathcal{F})$ will leave at least one of these paths between x and z , intact. Therefore x and z must lie in the same connected component in $G_{Z,W}(\mathcal{F})$. But $z \in V(\mathcal{B})$ and hence z must lie in C_1 and we assumed that v lies in C_2 . Contradiction. So we conclude that $G_{Z,W}(\mathcal{F})$ cannot have

more than one connected component. That is, removing any $d - k - 2$ vertices does not disconnect $G_Z(\mathcal{F})$. Hence $G_Z(\mathcal{F})$ is $(d - k - 1)$ -connected. $\bigcirc \bigcirc$

Theorem 1 shows that if we remove the vertices of a k -face from the graph of a d -polytope the remaining subgraph is *at least* $(d - k - 1)$ -connected. The following construction shows that this lower bound is tight when $k < d - 1$. (We do not use a d -simplex instead of the following construction, because removing the vertices of a k -face leaves only $d - k$ vertices in the graph of the simplex and it would not be meaningful to consider the $(d - k)$ -connectedness of the remaining subgraph).

Construction 1 : Let P be a simple d -polytope and v a vertex of P . Obtain Q by truncating the vertex v from P . That is, if H is a hyperplane such that $v \in H^+$ and $V(P) \setminus \{v\} \subset H^-$ (H^+ and H^- are the two open half-spaces determined by H) then

$$Q = (H \cup H^-) \cap P.$$

Q is a simple d -polytope. $F = H \cap Q$ is a facet of Q . More importantly, F is a $(d - 1)$ -simplex. Consider any k -face Z of F , $0 \leq k < d - 1$. Since Z is a k -simplex it has $k + 1$ vertices. Since $k < d - 1$ there is a vertex $z \in V(F) \setminus V(Z)$. z has d neighbours in Q and the vertices of Z are $k + 1$ of them. So in the subgraph induced by the vertex set $V(Q) \setminus V(Z)$, z has $d - k - 1$ neighbours and hence the subgraph $G(Q) \setminus G(Z)$ cannot be $(d - k)$ -connected. \bigcirc

From result 2.1 we know that the set of all vertices that are not in a given facet induce a subgraph that is *at least* 1-connected. We conclude by showing that this lower bound is tight as well.

Construction 2 : Let X be any $(d - 1)$ -polytope contained in a hyperplane H in R^d . Choose $v_1, v_2 \in R^d \cap H^+$ such that v_1 and v_2 are vertices of $Y = \text{conv}(\{v_1, v_2\} \cup X)$. Now choose a $v_3 \in H^+ \setminus Y$ such that $\text{conv}(v_1, v_3) \cap \text{int}(Y) \neq \emptyset$.

It is easy to see that v_1, v_2 and v_3 are vertices of $Q = \text{conv}(\{v_3\} \cup Y)$. Moreover we chose v_3 such that (v_1, v_3) is not an edge of Q . X is a facet of Q and the complement of $G(X)$ (i.e. $G(Q) \setminus G(X)$) is the path $v_1 \leftrightarrow v_2 \leftrightarrow v_3$ which is *not* 2-connected. \bigcirc

3 Acknowledgment

My advisor, Richard Pollack cleaned up the initial version to an extent where he practically rewrote the whole paper. It is with regret that I merely thank him for his patience, support and for all the discussions with him that led to this paper. The discussions with my advisor Richard Cole, Louis Billera and Victor Klee (about a conjecture of Micha Perles) were also very helpful.

References

- [1] M.L. Balinski. *On the graph structure of convex polyhedra in n -space*; Pacific Jour. Math. **11** (1961), 431-434.
- [2] G.T. Sallee. *Incidence graphs of convex polytopes*; Jour. Comb. Theory **2**, 466-506, 1967.
- [3] A. Brøndsted. *An introduction to convex polytopes*; Springer-Verlag, 1983.

- [4] P. McMullen and G.C. Shephard. *Convex polytopes and the upper bound conjecture*; London Math. Soc. Lecture Notes, vol. 3, London: Cambridge Univ. Press, 1971.
- [5] B. Grünbaum and T.S. Motzkin. *On polyhedral graphs*; In *Convexity* (V. Klee, ed.), Amer. Math. Soc. Proc. Symp. Pure Math. **7**, 285-290, 1963.
- [6] V. Klee. *A property of d -polyhedral graphs*; Jour. Math. and Mech., Vol. 13, No. 6, 1964.
- [7] V. Klee and P. Kleinschmidt. *Polytopal complexes and their relatives*; IMA preprint series # 384, Jan. 1988.

